

THE INTERACTION BETWEEN IRON, PHYTOPLANKTON AND THE CO₂ SYSTEM

Frank J. Millero
MAC/RSMAS/University of Miami
4600 Rickenbacker Causeway
Miami, FL 33149-1098
Tel: 305/361-4707
Fax: 305/361-4144
Internet: fmillero@rsmas.miami.edu
Award: N000149610927

LONG TERM GOALS

Our proposed work on the physical chemistry of coastal waters during the last cycle was not supported by ONR. We were given funds to support the wrap up of our past studies on the interaction of iron, phytoplankton and the CO₂ system. This report thus discusses our work over the last year on the iron work. It also discusses some of our measurements that were continued in coastal waters over the last year. This includes the development of flowing systems that can make continuous measurements of iron and nutrients in coastal waters. The basic aim of this work is to obtain an understanding of the chemical interactions between iron, phytoplankton, and the carbon dioxide system in the oceans. We hope to be able to determine how the growth of phytoplankton and its limitations on iron in some regions of the oceans affects the carbonate system. We also hope to determine how the speciation of iron affects its uptake by phytoplankton and examine how the competition of surface sites on phytoplankton with ligands in seawater affect the uptake.

OBJECTIVES

Iron is a biologically essential trace metal that because of its low levels may be limiting to the growth of phytoplankton in the oceans especially in areas of high nutrients and low chlorophyll. In addition iron may also affect the growth of phytoplankton in areas thought to be nitrogen limited and be a selective factor underlying the succession and diversity of phytoplankton species. It is now understood that the speciation of metals especially with organic ligands can play an important role in the availability and toxicity of metals in seawater. The inorganic and organic speciation of iron in seawater is not completely understood. Iron can exist in two oxidation states that have differences. Fe(II) is thought to be unstable in seawater and to be mostly in the free form. Fe(III) forms strong interactions with OH⁻ and is not very soluble in seawater. The possible existence of iron colloids adds a further complication to the activity of iron in the oceans. In this study we are attempting to understand the chemical interactions between iron, phytoplankton, and the carbon dioxide system in the oceans. We are making laboratory measurements and modeling studies to understand the speciation of iron in seawater and

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE The Interaction Between Iron, Phytoplankton and the CO2 System				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Miami,Rosensteil School of Marine and Atmospheric Science,4600 Rickenbacker Causeway,Miami,FL,33149				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

field measurements during the IRONEX studies to understand how the addition of iron to ocean waters can affect the carbonate system. This work is supported by the Office of Naval Research Physical Oceanography Program

APPROACH

Our laboratory measurements on the rates of oxidation and reduction of iron in natural waters are being made using the chemiluminescence techniques developed by O'Sullivan et al. (in press) and King et al. (1995). We also are examining the speciation of iron in seawater using polarography. We are modeling the speciation using Pitzer equations (Millero et al., 1995). Our field measurements have been made as part of the IRONEX I and II studies in the equatorial Pacific. Measurements were made of the changes in total carbon dioxide, total alkalinity and partial pressure of carbon dioxide in the iron enriched patch. More recent measurements have been made in Miami coastal waters.

WORK COMPLETED

Over the last year we have completed a number of studies on the rates of oxidation and reduction of iron in natural waters and the speciation of iron (see publications). We also have examined the speciation of iron in seawater using the Pitzer model. We have used this model to estimate the solubility of iron in seawater. As part of this work we have developed methods that can be used to make continuous measurements of pH, TA, TCO₂, pCO₂, NO₃, NO₂, PO₄, SiO₂ and Fe (see references). The overall results of the first IRONEX experiment have been published (Martin et al., 1994; Watson et al., 1994). We have also completed a data report which summarizes our CO₂ measurements and two papers (in press) to a special issue of Deep-Sea Research that will contain all the work resulting from the first experiment. Several papers were published this year on the effect of iron on carbon dioxide in the oceans (see references). We participated in the IRONEX II experiment May-June 1995 and made pH, TA, TCO₂ and pCO₂ measurements. Unlike the IRONEX I studies, iron was added more than once to the patch. This resulted in a larger pull-down of nutrients and carbon dioxide. The pCO₂ was decreased by 60 μ atm and the TCO₂ by 30 μ mol kg⁻¹. The TA showed little or no changes when corrected for salinity. The combined measurements of pCO₂ by our group and the nutrients by Sakamoto and Fredrich (IMBAR) show an excellent co-variation as found in the first experiments. More recently we have used two flowing systems to study coastal waters from Miami and through the Bahama Banks

RESULTS

The Ironex experiments have led to a better understanding of the effect of nutrients and the resultant grazing of zooplankton in natural systems. The issue of whether iron additions can sequester substantial quantities of atmospheric CO₂ will lead to further debates.

IMPACT

These experiments demonstrate that open ocean manipulative experiments are possible and ecological studies in the open ocean are no longer limited to passive observations. This will change significantly the way geochemical and ecological experiments are done in the future.

TRANSITIONS

Over the last year we have initiated studies of coastal waters near Miami in cooperation with Rod Zika's group here at the University of Miami.

RELATED PROJECTS

The carbon dioxide system in the Southern Ocean and the study of ionic interactions of Fe in coastal waters.

REFERENCES

- R.R. Bidigare, A. Fluegge, K. h. Freeman, K. l. Hanson, J.M. Hayes, D. Hollander, J.P. Jasper, L.L. King, E.A. Laws, J. Milder, F.J. Millero, R. Pancost. B.N. Popp, P.A. Steinberg and S.G. Wakeham, *Global Biogeochem. Cycles*, **11**(2), 279-292, 1997.
- D.W. King, H.A. Lounsbery and F.J. Millero, Rates and mechanism of Fe(II) oxidation at nanomolar concentrations, *Environ. Sci. Technol.*, **29**(3): 818-824, 1995.
- K. Lee, F.J. Millero, and R. Wanninkhof. The carbon dioxide system in the Atlantic Ocean, *J. Geophys. Res.*, in press.
- J. H. Martin, K. Coale, K.S. Johnson, S.E. Fitzwater, R.M. Gordon, S.J. Tanner, C.N. Hunter, V. Elrod, J Norwicki, T. Coley, R. Barber, S. Lindley, A. Watson, K. Van Scoy, C. Law, R. Ling, T. Stanton, J. Stockel, C. Collins, A. Anderson, R., M. Onrusek, M-Miguel Latasa, F.J. Millero, K. Lee, W. Yao, J.Z. Zhang, G. Friederich, C. Sakamoto, F. Chavez, K. Buck, S. Kolber, R. Green, P. Falkowski, S. W. Chisholm, F. Hoge, B. Swift, S. , S. Turner, P. Nightingale, P., Liss, and N. Tindale. The Iron Hypothesis: ecosystem test in Equatorial Pacific waters, *Nature*, **371**: 123-129, 1994.
- S. McElligott, R.H. Byrne, K. Lee, R. Wanninkhof, F.J. Millero, R.A.. Feely, Discrete water column measurements and thermodynamic calculations, *Mar. Chem.*, in press.
- F.J. Millero, W. Yao and J. Aicher, The speciation of Fe(II) and Fe(III) in seawater, *Mar. Chem.*, **50**, 21-39, 1995.
- F.J. Millero and R.N. Roy , A Chemical Equilibrium Model For The Carbonate System, in Natural Waters, *Croatia Chemica Acta*, **70**: 1-38, 1997.
- F.J. Millero. The effect of iron on carbon dioxide in the oceans. *Sci. Progr.* **80**(2), 147-168, 1997.
- F.J. Millero, W. Yao, K. Lee, J-Z. Zhang, and D.M. Campbell, Carbonate system in the waters near the Galapagos Islands, *Deep Sea Res.*, in press
- F.J. Millero, M. P. Roche and K. Lee, The total alkalinity of surface waters, *Mar. Chem.*, in press.

- F.J. Millero, The effect of ionic interactions on thermodynamic and kinetic processes in natural waters, NATO Book, in press.
- F.J. Millero, The influence of iron on carbon dioxide in surface seawater, NATO Book, in press.
- F.J. Millero, Solubility of Fe(III) in Seawater, *Earth Planet. Sci. Letters*, Submitted.
- D.W. O'Sullivan and F.J. Millero, Continuous measurement to the total inorganic carbon in surface waters, *Mar. Chem.*, in press.
- D. Pierrot, F.J. Millero, L.N. Roy, R.N. Roy, A. Doneski, and J. Nedersmidt, The Activity Coefficients of HCl in HCl-Na₂SO₄ Solutions from 0 to 50 °C and I = 6 m, *J. Solution Chem.*, 26(1): 31-45, 1997.
- M. Roche and F.J. Millero, Measurements of total alkalinity of surface waters using a continuous underway spectrophotometric technique, *Mar. Chem.*, in press.
- C. M. Sakamoto, F. J. Millero, W. Yao, G.E. Friederich, and F. P. Chavez, Surface seawater distributions of inorganic carbon and nutrients around the Galapagos Islands: Results from the PlumEx experiment using automated chemical mapping, *Deep Sea Res.*, in press.
- V.K. Sharma, J.O. Smith and F.J. Millero. Ferrate (VI) oxidation of hydrogen sulfide. *Environ Sci. Technol.*, 31: 2486-2491, 1997.
- P.A. Steinberg, F.J. Millero, and X. Zhu, Carbonate system response to iron enrichment, *Mar. Chem.*, submitted.
- A.J. Watson, C.S. Law, K. Van Scoy, M.I. F.J. Millero, W. Yao, G. Friederich, M.I. Liddicoat, R.H. Wanninkhof, R.T. Barber and K. Coale, Minimal effect of iron fertilization on sea-surface carbon dioxide concentrations., *Nature*, **371**: 143-145.
- X.R. Zhu, J.M. Prospero and F.J. Millero, Diel Variability of Soluble Fe(II) and Total Fe In Saharan Dust in the Trade Winds at Barbados, *J. Geophys. Res.*, 102,21,297-305.